

OPEN

Epidemiology Publish Ahead of Print

DOI: 10.1097/EDE.0000000000001137

Within-City Spatial Variations in Ambient Ultrafine Particle Concentrations and Incident Brain Tumors in Adults

Scott Weichenthal,^{*1,2} Toyib Olaniyan,³ Tanya Christidis,³ Eric Lavigne,² Marianne Hatzopoulou,⁴ Keith Van Ryswyk,² Michael Tjepkema,³ Rick Burnett⁵

¹ Department of Epidemiology, Biostatistics, and Occupational Health, McGill University, Montreal, Canada

² Air Health Science Division, Health Canada, Ottawa, Canada

³ Statistics Canada, Ottawa, Canada

⁴ University of Toronto, Department of Civil Engineering, Toronto, Ontario, Canada

⁵ Population Studies Division, Health Canada, Ottawa, Canada

*Corresponding Author: Dr. Scott Weichenthal, Department of Epidemiology, Biostatistics, and Occupational Health McGill University, 1020 avenue des Pins Ouest, Montreal, QC, Canada, H3A 1A2 Email: scott.weichenthal@mcgill.ca Tel: (514) 398-1584,

Running head: UFPs and Brain Tumors

Conflict of interest: None declared.

Financial Support: This study was funded by Health Canada. Dr. Weichenthal also received support from a GRePEC salary award funded by the Cancer Research Society, the Quebec Ministry of Economy, Science and Innovation, and FRQS (Fonds de Recherche du Québec-Santé).

Data Availability

All exposure data used in this manuscript are freely available. The CanCHEC cohort is not publicly available but can be accessed through Statistics Canada Research Data Centers conditional on the necessary approval/review procedures.

Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

ACCEPTED

Abstract

Background: Ambient ultrafine particles (UFPs, $<0.1 \mu\text{m}$) can reach the human brain but to our knowledge epidemiologic studies have yet to evaluate the relationship between UFPs and incident brain tumors.

Methods: We conducted a cohort study of within-city spatial variations in ambient UFPs across Montreal and Toronto, Canada among 1.9 million adults included in multiple cycles of the Canadian Census Health and Environment Cohorts (1991, 1996, 2001, and 2006). UFP exposures (3-year moving averages) were assigned to residential locations using land use regression models with exposures updated to account for residential mobility within and between cities. We followed cohort members for malignant brain tumors (ICD-10 codes C71.0-C71.9) between 2001 and 2016; Cox proportional hazards models (stratified by age, sex, immigration status, and census cycle) were used to estimate hazard ratios (HRs) adjusting for fine particle mass concentrations ($\text{PM}_{2.5}$), nitrogen dioxide (NO_2), and various sociodemographic factors.

Results: In total, we identified 1400 incident brain tumors during the follow-up period. Each $10,000/\text{cm}^3$ increase in UFPs was positively associated with brain tumor incidence (HR=1.112, 95% CI: 1.042, 1.188) after adjusting for $\text{PM}_{2.5}$, NO_2 , and sociodemographic factors. Applying an indirect adjustment for cigarette smoking and body mass index strengthened this relationship (HR=1.133, 95% CI: 1.032, 1.245). $\text{PM}_{2.5}$ and NO_2 were not associated with an increased incidence of brain tumors.

Conclusions: Ambient UFPs may represent a previously unrecognized risk factor for incident brain tumors in adults. Future studies should aim to replicate these results given the high prevalence of UFP exposures in urban areas.

Keywords: ultrafine particles, brain tumors, cohort study

Introduction

Outdoor air pollution and diesel exhaust are each classified as known human carcinogens based primarily on evidence related to lung cancer.^{1,2} In urban areas, diesel exhaust and other combustion processes are important sources of ambient ultrafine particles (<0.1 μm , UFPs)^{3,4} but little is known about the long-term health effects of these pollutants. To date, only a small number of studies, to our knowledge, have examined the relationship between UFPs and cancer including studies of lung⁵, prostate⁶, and postmenopausal breast cancer.⁷ In general, evidence from these studies is inconclusive and our knowledge remains limited with respect to the impact of ambient UFP exposures on cancer incidence in exposed populations.

Over the past several years, numerous studies have documented the ability of small inhaled particles to reach the brain.⁸ In particular, animal evidence suggests that a fraction of inhaled nanoparticles translocate to the brain when deposited in either the nasal epithelium (via the olfactory nerve) or the alveolar epithelium by entering into the systemic circulation and eventually crossing the blood brain barrier.⁸⁻¹² The rate of particle translocation generally depends on particle size and surface properties (e.g. charge, protein/lipid coatings) with smaller size favoring more rapid translocation.^{8,11} Moreover, the detrimental impacts of particle translocation may occur through particle retention at the target site (i.e. through the generation of reactive oxygen species in the brain¹²) or indirectly through inflammatory mediators produced in response to the particles.⁸ In general, while the rate of translocation of particles to the brain is slow, elimination rates are also low and thus particles may accumulate over time with long-term exposures.⁸

Importantly, the translocation of airborne particles to the brain is also supported by human evidence as Maher et al.¹³ recently reported the presence of combustion-related nanoparticles in the brains of subjects who lived in Mexico and the United Kingdom. In general, however, epidemiologic evidence related to outdoor air pollution and brain tumors is limited and somewhat inconsistent with four studies reporting positive associations¹⁴⁻¹⁷ and two others reporting no relationship.¹⁸⁻¹⁹ In the most recent study of European cohorts,¹⁷ PM_{2.5} absorbance (a marker for traffic-related air pollution) was most strongly associated with malignant brain tumors suggesting that traffic pollution may be an important source of exposure for these types of cancer. However, there have been no studies to date examining the relationship between brain tumors and ambient UFPs and these pollutants are of particular interest as they are known to reach the human brain. Here we address this important knowledge gap in a study of ambient UFPs and brain tumor incidence across Canada's two largest cities.

Methods

Study Population

Our study population was based on four cycles (1991, 1996, 2001, and 2006) of the Canadian Census Health and Environment Cohort (CanCHEC). These datasets were formed by linking Census long-form questionnaires (which collect data on approximately 20% of Canadian households every 5 years) to postal code histories and mortality and cancer incidence records through the Statistics Canada Social Linkage Data Environment.²⁰⁻²² The in-scope study population included non-institutionalized respondents under the age of 90 who lived in Toronto or Montreal for at least 1 year between 1998 and the end of follow-up in 2016. We used residential postal code histories (from annual income tax filings) to estimate time-varying UFP exposures to account for residential mobility both within and between cities. The CanCHEC

dataset was created under the authority of the Statistics Act and approved by the Executive Management Board (reference number: 045-2015) at Statistics Canada. This approval is equivalent to that of standard research ethics boards.

Brain Tumor Case Identification

We identified incident brain tumor cases (Malignant Neoplasms of the Brain: ICD-10 codes C71.0-C71.9) between 2001 and 2016 using the Canadian Cancer Registry. The follow-up period (i.e. the at-risk period) was restricted to years between 2001 and 2016 to avoid extrapolating UFP exposures many years in the past (i.e. for cohort members enrolled in 1991 and 1996). As a sensitivity analysis, outcome definition was limited to ICD-10 code C71.9 (malignant neoplasms of the brain, unspecified site) which captures gliomas (the most common type of malignant brain tumors in adults) and other brain tumor types with site unspecified within the brain. Other outcomes (including other types of cancer) were not examined.

Air Pollution Exposure Assessment

UFP exposures were assigned to residential locations (i.e. six digit postal codes, about the size of city block face or large apartment complex) using land use regression models for Montreal (2011-2012)⁴ and Toronto (2010-2011).²³ These models were developed using mobile monitoring data collected during the summer and winter months and explained the majority of spatial variations in ambient UFPs across each city (Montreal: $R_{CV}^2=0.60$; Toronto: $R_{CV}^2=0.50$). All UFP exposures were assigned as 3-year moving averages with a 1-year lag in order to account for residential mobility within or between cities. At the start of follow-up, each participant was assigned an exposure based on their residential location (from the land-use regression models) for the years 1998, 1999, and 2000. Each year, we updated this exposure to account for residential mobility. For example, exposure estimates for 2005 were based on the

average value for residential locations in 2002, 2003, and 2004. We did not project spatial variations in outdoor UFP concentrations back in time owing to the absence of long-term monitoring data for UFPs and a low correlation with other monitored pollutants.

We assigned long-term estimates (i.e. 3-year moving averages with 1-year lag) of ambient fine particulate air pollution (PM_{2.5}) and nitrogen dioxide (NO₂) concentrations to cohort members as previously described.²⁴⁻²⁵ Annual average concentrations were available for PM_{2.5} exposures assigned during follow-up and long-term estimates of spatial variations in NO₂ concentrations reflected the year 2006. Briefly, the PM_{2.5} exposure model was based on a combination of aerosol optical depth, chemical transport model, and land use information and estimated concentrations at a 1-km² spatial resolution.²⁴ Ambient NO₂ concentrations were estimated using a national land use regression model including parameters such as satellite-derived NO₂ and distances to major roads and highways. The primary purpose of including long-term estimates of ambient PM_{2.5} and NO₂ concentrations was to evaluate the sensitivity of UFP/brain tumor associations to spatial variations in other ambient air pollutants.

Statistical Analysis

Cox proportional hazard models²⁶ were used to estimate the association between ambient UFP concentrations and brain tumor incidence in Montreal and Toronto. A directed acyclic graph (DAG) outlining the conceptual relationship between UFPs and brain tumors (and potential confounding factors) is provided in eFigure 1; <http://links.lww.com/EDE/B613>. We examined several models to explore this relationship. First, we developed a base model including UFP concentrations and strata for age (5-year groups), sex, immigration status, and census year (i.e. 1991, 1996, 2001, and 2006). We used strata variables to allow for different baseline hazard functions across age, sex, immigrant status, and census year. Next, we examined the impact of

adding covariates for SES variables (described in Table 1) (visible minority status, occupational level, educational attainment, marital status, and income quintile), PM_{2.5}, and NO₂ both individually and combined. In constructing the visible minority variable, we grouped white and aboriginal cohort members together as aboriginal people made up a small portion of overall participants (11,300 of 1.9 million, or 0.6%) and person–years (116,100 of 25,707,900 person–years, or 0.45%). All UFP hazard ratios are scaled per 10,000 particles/cm³. All descriptive statistics were rounded to the nearest hundred for confidentiality (as required by Statistics Canada).

Indirect Adjustment for Smoking and Body Mass Index

As sensitivity analyses, we applied a method of indirect adjustment for smoking and body mass index as these unmeasured parameters have been associated with brain tumors in previous studies.²⁷⁻²⁸ As indicated in the DAG presented in eFigure 1; <http://links.lww.com/EDE/B613>, smoking and body mass index are not causes of long-term outdoor UFP concentrations (i.e. intervening on individual-level smoking or BMI is not expected to change long-term outdoor UFP concentrations) and thus are not represented in the DAG as confounders in the typical sense. Nevertheless, chance associations could confound the relationship between outdoor UFP concentrations and brain tumor incidence, and the indirect adjustment method was applied to address this possibility. Likewise, SES variables (i.e. income, education) could be related to smoking/BMI and this relationship could in turn bias the observed relationship between outdoor UFP concentrations and brain tumors. Applying the indirect adjustment method for smoking and BMI also addresses this possibility.

A detailed description of the indirect adjustment method is available elsewhere.²⁹ Briefly, this method requires information on the correlation between observed and missing risk factors from an ancillary data source (e.g. national health surveys) as well as estimates of the relationship between the missing risk factors and brain tumor incidence. For our analyses, information related to the correlation between observed and missing risk factors was obtained from multiple cycles of Canadian Community Health Survey (CCHS) (2001, 2003, 2005, 2007, 2009, 2011, 2013); the CCHS is a bi-annual national health survey that collects information on the health status and behaviours (i.e. smoking and BMI) of non-institutionalized Canadians aged 12 years and older. Risk estimates for the relationship between brain tumors and smoking and body mass index were obtained from the published literature.²⁷⁻²⁸ The CCHS and CanCHEC databases were compared with respect to UFP concentrations across age, sex, and SES variables; median differences were small in magnitude (typically less than 1000/cm³) supporting the use of the CCHS database in estimating correlations between observed and missing risk factors (eTable 1; <http://links.lww.com/EDE/B613>).

Concentration Response Relationship

The shape of the concentration–response relationship between outdoor UFP concentrations and brain tumor incidence was examined using the Shape Constrained Health Impact Function (SCHIF) method developed by Nasari et al.³⁰ Specifically, this method utilizes flexible concentration transformations (i.e. linear, sub-linear, supra-linear, and sigmodal) within a Cox model and produces biologically plausible concentration–response relationships that increase monotonically with concentration. The final concentration-response curve was based on weights from the three best fitting models identified by the SCHIF method.

Results

Cohort characteristics are presented in Table 1. In total, we followed 1,938,100 adults (25,707,900 person–years) in Montreal and Toronto between 2001 and 2016 and 1400 incident cases of brain tumors were included in the analyses. We excluded approximately 10 million person–years based on our inclusion criteria (i.e. less than 90 years of age at start of follow-up and resided in Toronto or Montreal for at least 1 year between 1996 and 2016). Incident brain tumors were more common among women (n=800) than men (n=600) and were primarily diagnosed between age 40-69 years (eTable 2; <http://links.lww.com/EDE/B613>).

Mean UFP values assigned to all in-scope person–years (i.e. 3-year moving averages with a 1-year lag) were 26,697/cm³ (SD= 8030/cm³) and ranged from 6697/cm³ to 97,158/cm³. Spatial variations in ambient UFPs were weakly correlated with PM_{2.5} (r= 0.14) and NO₂ (r= 0.20) whereas PM_{2.5} and NO₂ were more strongly correlated (r=0.66). The low correlation between UFPs and other air pollutants is commonly observed in Canada;³¹ this relates in part to the fact that the prevalence of diesel vehicles (a major source of UFPs) is low and thus it is possible to have areas with elevated NO₂ concentrations (from gasoline vehicles) without major elevations in UFPs. Moreover, PM_{2.5} tends to be a regional pollutant without major spatial variations within-cities. The mean PM_{2.5} concentration in this study was 8.68 µg/m³ (SD=2.42 µg/m³) and the mean NO₂ concentrations was 20.2 ppb (SD= 8.43 ppb).

Hazard ratios describing the relationship between ambient UFPs and incident brain tumors are listed in Table 2. UFPs were consistently associated with increased brain tumor incidence in all models examined and the hazard ratio remained elevated after adjustment for socio-demographic factors as well as PM_{2.5} and NO₂ (HR=1.112, 95% CI: 1.042, 1.188). A

slightly stronger association was observed in the fully adjusted model (i.e. all SES variables, PM_{2.5}, and NO₂) when the outcome was limited to ICD-10 code C71.9 (HR=1.170, 95% CI: 1.000, 1.372); however, this estimate was less precise owing to a smaller number of cases (n=200 for ICD-10 code C71.9).

Applying the indirect adjustment for cigarette smoking and body mass index (with adjustment for all SES variables, PM_{2.5}, and NO₂) slightly increased the strength of the observed association between UFPs and brain tumors (HR=1.133, 95%CI: 1.032, 1.245). This association was stronger in Toronto (HR=1.171, 95% CI: 1.021, 1.343) than Montreal (HR=1.022, 95%CI: 0.814, 1.283) but city-specific risk estimates were less precise and more than half (n=850) of the cases were from Toronto. The concentration–response curve for ambient UFP concentrations and incident brain tumors is shown in Figure 1. This relationship was s-shaped with a smaller slope at lower concentrations.

Spatial variations in PM_{2.5} and NO₂ were not positively associated with brain tumor incidence (eTables 3-4; <http://links.lww.com/EDE/B613>). Specifically, hazard ratios for interquartile changes in PM_{2.5} (per 3 µg/m³) and NO₂ (per 10 ppb) in fully adjusted models including all SES variables and UFPs were 0.907 (95% CI: 0.762, 1.079) and 0.950 (95% CI: 0.851, 1.061), respectively.

Discussion

We conducted a population-based study of within-city spatial variations in ambient UFP concentrations and incident brain tumors in Canada's two largest cities. In general, our findings suggest that ambient UFPs are positively associated with brain tumor incidence and these pollutants may represent a previously unrecognized risk factor for brain tumors. Moreover, the relationship between UFPs and brain tumors was not explained by other air pollutants such as

PM_{2.5} mass concentrations or NO₂ which are more frequently monitored in urban environments. On an absolute scale, the magnitude of the observed association suggests that each 10,000/cm³ increase in 3-year average ambient UFP concentration contributes to approximately 1 new case per 100,000 population (assuming a baseline age-standardized incidence rate of 8 per 100,000).³²

Existing evidence related to outdoor air pollution and brain tumors is inconsistent. For example, McKean-Cowdin et al.¹⁸ examined the relationship between PM_{2.5} and NO₂ and brain tumor mortality in the Cancer Prevention Study-II cohort (1982-2000) and reported small inverse associations similar in magnitude to those observed in the present study. Raaschou-Nielsen¹⁵ reported a positive association between NO_x (a marker for traffic pollution) and incident brain tumors in the Danish Diet, Cancer, and Health cohort in 2011 but a weaker association was observed in a follow-up case-control study including a much larger number of cases.¹⁹ Similarly, weak positive associations were observed between brain tumors and PM_{2.5}, NO₂, and NO_x in the Danish Nurse Cohort with stronger associations observed among obese subjects and those with lower levels of physical activity.¹⁶ Finally, the most recent study of air pollution and brain tumors reported weak positive associations between PM_{2.5} and NO₂ and brain tumors across multiple European cohorts with the strongest association observed for PM_{2.5} absorbance (a marker of traffic pollution).¹⁷ Collectively, existing evidence suggests that traffic-related air pollution may be associated with a small increased risk of brain tumor incidence/mortality but few studies have examined this relationship to date. Moving forward, particular attention should be paid to UFPs as they are known to reach the brain⁸⁻¹³ and our findings suggest a positive association with brain tumors independent of PM_{2.5} or NO₂ (which were not positively associated with brain tumor incidence). Moreover, most brain tumors are rapidly fatal and few known risk

factors have been identified for these types of cancer; therefore, it is important to identify potential environmental causes of brain tumors to inform future policy interventions.

This study had a number of important advantages including high-resolution estimates of within-city spatial variations in ambient UFP concentrations, updated exposure information (i.e. UFPs, NO₂, and PM_{2.5}) for subjects moving both within or between-cities, a large study population, incident cases, and detailed individual-level information on a number of important socio-economic factors. However, it is also important to note several limitations. First, the exposure models used to estimate spatial variations in outdoor UFP concentrations in Montreal (2011-2012) and Toronto (2010-2011) were based on data collected toward the end of the follow-up period (2001-2016) and exposure measurement error almost certainly impacted our results. This error likely has a temporal component (impacted by changes in emissions over time) and a spatial component (impacted by local changes in infrastructure that could influence the movement of traffic sources through space). While this error likely contributed to imprecision in our results, a systematic difference in the magnitude of exposure error by case status seems unlikely and the overall impact of exposure measurement error was likely a bias toward the null. Another possible concern is that trends in brain cancer incidence may influence our results (for example, if incidence was higher/lower closer to the time of land-use regression model development this could contribute to differential exposure measurement error); however, data compiled by the Canadian Cancer Society indicate that age-standardized incidence rates for brain/CNS tumors were been stable between 2001-2010.³³ Nevertheless, since outdoor UFP concentrations have likely decreased over time (owing to improved vehicle efficiency), our data may not be appropriate in identifying absolute threshold concentrations for UFP impacts on brain tumors if overall exposure levels were elevated toward the beginning of the follow-up period

(data are not currently available to verify or refute this hypothesis). Furthermore, as for UFPs, exposure measurement error likely also impacted our estimates of PM_{2.5} and NO₂ and this may have resulted in residual confounding by these factors. However, including PM_{2.5} and/or NO₂ in the models tended to increase hazard ratios for UFPs and brain tumors and thus more precise estimates would be expected to further strengthen this relationship. As in all studies, we also cannot rule out residual confounding of the UFP–brain tumor relationship by some unmeasured factors; however, it is unclear what these factors may be as there are few known risk factors for brain tumors and to explain our results these factors would also have to be related to spatial variations in outdoor UFPs.

A second limitation was the absence of individual level smoking and body mass index data for CanCHEC cohort members. As outlined in the DAG presented in the eFigure 1; <http://links.lww.com/EDE/B613>, individual-level smoking and BMI are not likely causes of long-term exposures to outdoor UFPs (i.e. changing individual-level smoking status or BMI is not expected to change long-term outdoor UFP levels) and thus these parameters are not identified in the DAG as confounders in the typical sense. However, chance associations between smoking or BMI and UFPs could still confound the analysis; thus, we conducted an indirect adjustment for these unmeasured risk factors to evaluate the sensitivity of our findings to this potential source of bias. The results of the indirect adjustment for smoking and BMI suggested a stronger association between UFPs and brain tumors, thus suggesting that confounding by smoking or BMI is not a likely explanation for the observed relationship between UFPs and brain tumors. Similarly, we did not have individual-level information on other potential risk factors for brain tumors including family history or life-time exposure to ionization radiation. It was not possible to indirectly adjust for these factors and we cannot rule

out potential confounding (away from or toward the null) by these variables if by chance they were systematically related to outdoor UFP concentrations across Montreal and Toronto.

Finally, given the rarity of brain tumors, our analyses focussed primarily on total malignant brain tumors (ICD-10 codes C71.0-C71.9: Malignant Neoplasms of the Brain) to maximize precision and does not provide an in-depth examination of specific tumor sub-types. As a result, if UFPs are only associated with specific types of brain tumors, the hazard ratios presented above may underestimate the true impact of UFPs on these specific tumor sub-types.

In conclusion, we conducted to our knowledge the first cohort study of outdoor UFP concentrations and incident brain tumors and noted a consistent positive association. This relationship was robust to adjustment for various sociodemographic factors as well as indirect adjustment for smoking and body mass index. Future studies should aim to replicate our findings as UFPs are known to reach the human brain and exposure prevalence is high in urban areas around the world.

References

1. Benbrahim-Tallaa L, Baan RA, Grosse Y, et al. Carcinogenicity of diesel-engine and gasoline-engine exhausts and some nitroarenes. *Lancet Oncology* 2012; 13: 663-664.
2. Loomis D, Grosse Y, Lauby-Secretan B, et al. The carcinogenicity of outdoor air pollution. *Lancet Oncology* 2013; 14: 1262-1263.
3. Weichenthal S, Van Ryswyk K, Kulka R, Sun L, Wallace L, Joseph L. In-vehicle exposure to particulate air pollution in Canadian metropolitan areas: the urban transportation exposure study. *Environ Sci Technol* 2015; 49: 597-605.
4. Weichenthal S, Van Ryswyk K, Goldstein A, Bagg S, Shekharizfard M, Hatzopoulou M. A land use regression model for ambient ultrafine particles in Montreal, Canada: a comparison of linear regression and a machine learning approach. *Environ Res* 2016; 146: 65-72.
5. Weichenthal S, Bai L, Hatzopoulou M, Van Ryswyk K, et al. Long-term exposure to ambient ultrafine particles and respiratory disease incidence in Toronto, Canada: a cohort study. *Environ Health* 2017; 16: 64.
6. Weichenthal S, Lavigne E, Valois MF, et al. Spatial variations in ambient ultrafine particle concentrations and the risk of incident prostate cancer: a case-control study. *Environ Res* 2017; 156: 374-380.
7. Goldberg MS, Labreche F, Weichenthal S, et al. The association between the incidence of postmenopausal breast cancer and concentrations at street-level of nitrogen dioxide and ultrafine particles. *Environ Res* 2017; 58: 7-15.
8. Heusinkveld HJ, Wahle T, Campbell A, et al. Neurodegenerative and neurological disorders by small inhaled particles. *Neurotoxicology* 2016; 56: 94-106.

9. Mitchell R, Maher BA. Evaluation and application of biomagnetic monitoring of traffic-derived particulate pollution. *Atmos Environ* 2009; 43: 2095-2103.
10. Kreyling WG. Discovery of unique and ENM-specific pathophysiologic pathways: comparison of the translocation of inhaled iridium nanoparticles from nasal epithelium versus alveolar epithelium towards the brain of rats. *Toxicol Appl Pharmacol* 2016; 299: 41-46
11. Oberdorster G, Elder A, Rinderknecht A. Nanoparticles and the brain: cause for concern? *J Nanosci Nanotechnol* 2009; 9: 4996-5007.
12. Oppenheim HA, Lucero J, Guyot AC, et al. Exposure to vehicle emissions results in altered blood brain barrier permeability and expression of matrix metalloproteinases and tight junction proteins in mice. *Part Fibre Toxicol* 2013; 10: 62.
13. Maher BA, Ahmed IAM, Karloukovski V, et al. Magnetite pollution nanoparticles in the human brain. *PNAS* 2016; 113: 10797-10801.
14. Boeglin ML, Wessels D, Henshel D. An investigation of the relationship between air emissions of volatile organic compounds and the incidence of cancer in Indiana counties. *Environ Res* 2006; 100: 242-254.
15. Raaschou-Nielsen O, Andersen ZJ, Hvidberg M, et al. Air pollution from traffic and cancer incidence : a Danish cohort study. *Environ Health*. 2011;10: 67.
16. Jørgensen JT, Johansen MS, Ravnskjaer L, et al. Long-term exposure to ambient air pollution and incidence of brain tumours: The Danish Nurse Cohort. *Neurotoxicology*. 2016; 55:122-130.
17. Andersen ZJ, Pedersen M, Weinmayr G, et al. Long-term exposure to ambient air pollution and incidence of brain tumour: the European Study of Cohorts for Air Pollution Effects (ESCAPE). *Neuro Oncol* 2018; 20: 420-432.

18. McKean-Cowdin R, Calle EE, Peters JM, et al. Ambient air pollution and brain cancer mortality. *Cancer Causes Control* 2009; 20: 1645-1651.
19. Poulsen AH, Sørensen M, Andersen ZJ, Ketzel M, Raaschou-Nielsen O. Air pollution from traffic and risk for brain tumors: a nationwide study in Denmark. *Cancer Causes Control*. 2016; 27:473-480.
20. Christidis T, Labrecque-Synnott F, Pinault I, Saidi A, Tjepkema M. 2018. The 1996 CanCHEC: Canadian Census Health and Environment Cohort profile. *Analy Studies: Methods Ref*. Statistics Canada No. 11-633-X no. 013.
21. Pinault L, Fines P, Labrecque-Synnott F, Saidi A, Tjepkema M. 2016. The Canadian Census-Tax-Mortality Cohort: A 10-Year Follow-Up. *Anal Studies: Methods Ref*. Statistics Canada No. 11-633-X no. 003.
22. Wilkins R, Tjepkema M, Mustard C, Choinière R. 2008. The Canadian census mortality follow-up study, 1991 through 2001. *Health Rep* 19:25–43. Statistics Canada No. 82-003-XPE.
23. Weichenthal S, Van Ryswyk K, Goldstein A, et al. Characterizing the spatial distribution of ambient ultrafine particles in Toronto, Canada: a land use regression model. *Environ Poll* 2016; 208: 241.
24. van Donkelaar A, Martin RV, Spurr RJD, Burnett RT. High resolution satellite-derived PM_{2.5} for optical estimation and geographically-weighted regression over North America. *Environ Sci Technol* 2015; 49: 10482-10491.
25. Hystad P, Setton E, Cervantes A, et al. Creating national air pollution models for population exposure assessment in Canada. *Environ Health Perspect* 2011; 119:1123-1129.
26. Cox DR. 1972. Regression models and life tables. *J R Stat Soc B* 20:187-220.

27. Li HX, Peng XX, Zong Q, et al. Cigarette smoking and risk of adult glioma: a meta-analysis of 24 observational studies involving more than 2.3 million individuals. *OncoTargets Ther* 2016; 9: 3511–3523.
28. Niedermaier T, Behrens G, Schmid D, Schlecht I, Fischer B, Leitzmann MF. Body mass index, physical activity, and risk of adult meningioma and glioma. *Neurology Oct* 2015; 85: 1342-1350.
29. Shin HH, Cakmak S, Brion O, et al. Indirect adjustment for multiple missing variables applicable to environmental epidemiology. *Environ Res* 2014; 482-487.
30. Nasari MM, Szyszkowicz M, Chen H, et al. A class of non-linear exposure-response models suitable for health impact assessment applicable to large cohort studies of ambient air pollution. *Air Qual Atmos Health* 2016; 9:961-972.
31. Weichenthal S, Hatzopoulou M, Goldberg M. Exposure to traffic-related air pollution during physical activity and acute changes in blood pressure, autonomic and microvascular function in women: a cross-over study. *Part Fibre Toxicol* 2014; 11: 70.
32. Canadian Cancer Statistics Advisory Committee. Canadian Cancer Statistics 2018. Toronto, ON: Canadian Cancer Society; 2018. Available at: cancer.ca/Canadian-Cancer-Statistics-2018-EN (accessed [April 4, 2019]).
33. Canadian Cancer Society's Advisory Committee on Cancer Statistics. Canadian Cancer Statistics 2015. Toronto, ON: Canadian Cancer Society; 2015. Available: <http://www.cancer.ca/~media/cancer.ca/CW/cancer%20information/cancer%20101/Canadian%20cancer%20statistics/Canadian-Cancer-Statistics-2015-EN.pdf?la=en>

FIGURES

Figure 1. Concentration–response relationship between ambient UFP concentrations and brain tumour incidence in Montreal and Toronto, Canada (2001-2016). The model is stratified by 5-year age groups, sex, census cycle (1991, 1996, 2001, 2006), and immigrant status and adjusted for visible minority status, occupational level, education, marital status, and income quintile. Dashed red lines indicated the 95% confidence interval. Hazard ratios (the solid black line) for the concentration–response curve are calculated in reference to the 1st percentile of UFP concentrations in the cohort ($15,767/\text{cm}^3$) up to the 99th percentile ($56,767/\text{cm}^3$).

ACCEPTED

TABLES

Table 1. Descriptive statistics for CanCHEC cohort members (1991, 1996, 2001, 2006)

Covariates	Person-years ^a	UFP (counts/cm ³)	
		Mean	SD
All	25,707,900	26,697	8030
Sex			
Male	13,818,800	26,722	8015
Female	11,889,100	26,668	8048
Age at enrollment (years)			
25 to 29	2,543,400	26,639	7958
30 to 39	6,284,700	26,607	8010
40 to 49	6,255,100	26,532	7958
50 to 59	4,662,400	26,606	8003
60 to 69	3,432,800	26,971	8150
70 to 79	2,022,100	27,182	8227
80 to 89	507,800	27,207	8025
Visible Minority Status			
White or Aboriginal ^b	19,065,900	26,483	7819
Visible minority	6,642,000	27,313	8577
Immigrant status			
Non-immigrant	13,697,700	25,923	7466
immigrant	12,010,200	27,580	8544
Marital status			

Single	5,137,100	26,372	7621
Common-law	2,090,800	25,307	6708
Married	14,641,400	26,989	8318
Separated	750,100	27,190	8404
Divorced	1,722,700	26,376	7729
Widowed	1,366,000	27,056	8116
Educational attainment			
Not completed high school	6,532,100	27,150	8195
High school with/without trades certificate	7,759,100	26,818	8298
Post-secondary non-university	4,652,500	26,615	8134
University degree	6,764,400	26,177	7433
Occupational level			
Management	2,180,300	26,346	7744
Professional	4,100,900	25,962	7293
Skilled, technical & supervisory	4,719,000	26,650	8089
Semi-skilled	5,560,300	26,994	8407
Unskilled	1,806,800	27,236	8382
Not applicable	7,340,700	26,885	8056
Income adequacy quintile			
1st quintile - lowest	5,588,900	26,913	8039
2nd quintile	5,290,900	27,009	8237
3rd quintile	5,009,000	26,812	8229

4th quintile	4,730,200	26,638	8128
5th quintile - highest	5,089,000	26,078	7458

^aNumbers were rounded to the nearest 100 for confidentiality; ^b White and aboriginal cohort members were grouped together as aboriginal people made up a small portion of overall participants (11,300 of 1.9 million, or 0.6%) and person–years (116,100 of 25,707,900 person–years, or 0.45%).

ACCEPTED

Table 2. Hazard ratios (95% confidence intervals) for UFP concentrations (per 10,000/cm³) and incident brain tumours (n=1400) in Montreal and Toronto, Canada (2001-2016)

Model	HR (95% CI)
Base Model ^a	1.099 (1.033, 1.169)
<i>Base Model + SES^b Variables (Individually)</i>	
+ Visible Minority Status	1.095 (1.029, 1.166)
+ Occupational Level	1.100 (1.034, 1.706)
+ Education	1.101 (1.035, 1.171)
+ Marital Status	1.097 (1.031, 1.167)
+ Income Quintile	1.099 (1.034, 1.170)
<i>Base Model + Co-Pollutants (Individually)</i>	
+ NO ₂	1.114 (1.045, 1.187)
+ PM _{2.5}	1.108 (1.039, 1.180)
Fully Adjusted Multi-Pollutant Models	
All SES	1.094 (1.028, 1.164)
All SES + NO ₂	1.104 (1.035, 1.177)
All SES + PM _{2.5}	1.104 (1.036, 1.177)
All SES + PM _{2.5} + NO ₂	1.112 (1.042, 1.188)
All SES + PM _{2.5} + NO ₂ + Indirect Adjustment for Smoking and BMI	1.133 (1.032, 1.245)

^a Base model is stratified by 5-year age groups, sex, census cycle (1991, 1996, 2001, 2006), and immigrant status. ^b SES, socio-economic status variables (visible minority status, occupational level, education, marital status, and income quintile). BMI indicates body mass index.

Figure 1

